

NOTES ON INTERSECTION COMPLEXES AND L -FUNCTIONS

JONATHAN WANG

ABSTRACT. This are notes from informal lectures on *Intersection complexes and unramified L -factors* (joint paper with Yiannis Sakellaridis). Most of the notes are concerned with background material not covered in the paper – I claim no originality in these parts (except for errors) and appropriate references are given.

CONTENTS

1. Integral representations of L -functions	2
1.1. Schwartz space	2
1.2. Sakellaridis' conjecture	2
1.3. Local conjecture of Sakellaridis–Venkatesh	3
1.4. Generalized Ichino-Ikeda conjecture	5
1.5. Understanding the Euler product	6
2. Examples	7
2.1. Classical examples	7
2.2. New examples	7
3. Summary of results	8
3.1. Previous work	8
3.2. Main result on Plancherel measure	9
3.3. Recipe for ρ_X	10
3.4. Assumption $\check{G}_X = \check{G}$	10
3.5. Bernstein asymptotics	10
3.6. Main result in terms of asymptotics	12
3.7. Main result in terms of Radon transform	12
3.8. Formula for Φ^0	13
4. Geometric techniques	13
4.1. IC function	13
4.2. Models for the formal arc space	14
4.3. Zastava space and the graded factorization property	15
4.4. Central fibers	16
4.5. Crystals	17
4.6. Further details	17
References	18

1. INTEGRAL REPRESENTATIONS OF L -FUNCTIONS

In this section we give an overview of the program that proposes to associate to any affine spherical variety an integral representation of a certain L -function.

1.1. **Schwartz space.** For now, let \mathbb{k} be any global field, and G a connected split¹ reductive group over \mathbb{k} . Let X be an affine spherical variety over \mathbb{k} . We assume:

- X has good integral models away from a finite set of places (including archimedean). If \mathbb{k} is a function field we just assume X is defined over \mathbb{F}_q . For number fields, such good integral models exist by [Sa12, Proposition 2.3.5].
- If \mathbb{k} is a function field, assume X behaves “like a characteristic 0” variety.

Let \mathbb{A} denote the adèles of \mathbb{k} . For a place v of \mathbb{k} , let F_v denote the local field with ring of integers \mathcal{O}_v . Let $[G] = G(\mathbb{k}) \backslash G(\mathbb{A})$.

1.1.1. Let $X^\bullet \subset X$ denote the open G -orbit². In [Sa12, §3], Sakellaridis conjectures that there exists a nice Schwartz space

$$\mathcal{S}(X(\mathbb{A})) := \bigotimes_v^I \mathcal{S}(X(F_v)) \subset C^\infty(X^\bullet(\mathbb{A}))$$

which is a restricted tensor product of local Schwartz spaces with respect to a certain *basic function* $\Phi_v^0 \in C^\infty(X^\bullet(F_v))$. The Schwartz space $\mathcal{S}(X(F_v))$ consists of functions on $X^\bullet(F_v)$, but its definition will depend on the affine embedding $X^\bullet \hookrightarrow X$. For example, the basis function Φ_v^0 should have support contained in $X(\mathcal{O}_v) \cap X^\bullet(F_v)$.

We will return to the basis function in more detail later.

1.2. **Sakellaridis’ conjecture.** Starting from a Schwartz function $\Phi \in \mathcal{S}(X(\mathbb{A}))$, we can form the X -theta series $\Theta_X(\Phi, g) \in C^\infty([G])$.

$$(1.1) \quad \Theta_X(\Phi, g) = \sum_{\gamma \in X^\bullet(\mathbb{k})} g \cdot \Phi(\gamma)$$

As long as Φ has some reasonable growth conditions, this sum will be absolutely convergent. In fact $\Theta_X(\Phi, g)$ will be of moderate growth in g (see [Sa12, Proposition 3.1.3], [BP, Proposition A.1.1(ix)]).

Remark 1.2.1. The formula (1.1) looks similar to the formula for the theta function in the theta correspondence: there we summed over E^* where E^* is a Lagrangian vector space in the $G = \mathrm{Sp}(E \oplus E^*)$ -Hamiltonian variety $T^*(E^*) = E \oplus E^*$. Here we are summing over X^\bullet , which is a Lagrangian variety in the G -Hamiltonian variety T^*X^\bullet . These two construction should fall within the same unifying framework of [BZSV]. However the G -action on $\mathcal{S}(E^*)$ is more complicated because E^* is not a G -stable Lagrangian while X^\bullet is. Slogan: it’s harder to quantize a G -Hamiltonian variety M if you don’t have a G -stable Lagrangian.

Definition 1.2.2. We define a “zeta integral”

$$\mathcal{Z}(\chi, \Phi, f) := \mathcal{P}_\Phi(\chi \cdot f) = \int_{[G]} \chi(g) \cdot \Theta_X(\Phi, g) \cdot f(g) dg$$

where χ is an idele class character of G and $f \in \mathcal{A}_0(G)$ is a cusp form.

¹There should be generalizations to non-split groups, but there may be subtleties with the Galois group.

² $X_{\mathbb{k}}$ has a unique $G_{\mathbb{k}}$ -orbit by sphericity, which descends to a unique G -orbit of X .

This zeta integral *a priori* converges absolutely when $\log \chi$ lies in a translate of a rational cone prescribed by X (these are called sufficiently X -positive characters). (Even though $\Theta_X(\Phi, g)$ has moderate growth, the integral may not converge when $Z(G)^0$ is non-trivial.)

Naively we would like to say that the “gcd” of $\mathcal{Z}(\chi, \Phi, f)$ over all Φ should give us an L -function. This is sometimes the case, but the general expectation is more nuanced – we refer to §1.4 for the precise expected relation to L -functions.

Remark 1.2.3. In order to replace χ with $s \in \mathbb{C}$ (for L -function comparisons) we would need to have a canonical family of idele class characters. This requires that $Z(G)^0$ be non-trivial. We can always reduce to this assumption when X is not homogeneous: We can replace G by $G \times^{Z(G)^0} Z(X^\bullet)$ so that $Z(G)^0 = Z(X^\bullet)$, where $Z(X^\bullet) := \text{Aut}^G(X^\bullet)^0$ and $\text{Aut}^G(X^\bullet) = N_G(H)/H$ is the group of G -automorphisms of $X^\bullet = H \backslash G$. It follows from some structural results on spherical varieties (cf. [Sa12, Proposition 2.2.6]) that if X^\bullet is quasi-affine but not affine, then $Z(X^\bullet)$ is a non-trivial torus. (It is a non-trivial theorem that $\text{Aut}^G(X^\bullet)$ is always diagonalizable.)

Conjecture 1.2.4 (Weak form, [Sa12, Conjecture 3.2.4]). *For any $\Phi \in \mathcal{S}(X(\mathbb{A}))$ and $f \in \mathcal{A}_{\text{cusp}}(G)$, the integral $\mathcal{Z}(\chi, \Phi, f)$ admits meromorphic continuation to the space of all idele class characters χ on G . One can further hope for a “Fourier transform” and functional equation.*

Remark 1.2.5. This conjecture is a generalization of the conjecture of Braverman–Kazhdan [BrK] seeking to generalize Godement–Jacquet theory. In the Braverman–Kazhdan conjecture, you start with some G with a “determinant” map $G \rightarrow \mathbb{G}_m$ and an irreducible representation ρ of \tilde{G} (satisfying some compatibility condition). Then to ρ you can attach a reductive monoid $M_\rho \supset G$, which is an affine spherical variety for $G \times G$. Then Braverman–Kazhdan’s Schwartz space $\mathcal{S}_\rho(G(\mathbb{A}))$ is just $\mathcal{S}(M_\rho(\mathbb{A}))$ in our notation, with the same conjectural properties. In that case we hope that the “gcd” of $\mathcal{Z}_{M_\rho}(|\det|^s, \Phi, f)$ equals the L -function associated to ρ .

There is also the following strong form of the conjecture, which seems far out of reach since it is not even known for parabolic Eisenstein series, i.e., the case $X = \overline{N_P} \backslash \overline{G}^{\text{aff}}$, where N_P is the unipotent radical of a parabolic P which is not a Borel.

Conjecture 1.2.6 (Strong form, [Sa12, Conjecture 3.2.2]). *For every $\Phi \in \mathcal{S}(X(\mathbb{A}))$, the X -Eisenstein series*

$$\mathcal{E}_X(\chi, \Phi, g) = \int_{Z(X^\bullet)(\mathbb{A})} \Theta_X(z \cdot \Phi, g) \chi(z) dz$$

originally defined for sufficiently X -positive idele class characters, admits meromorphic continuation everywhere.

For the problem of meromorphic continuation, it is not really important what the local Schwartz spaces $\mathcal{S}(X(F_v))$ are: the integral can have arbitrary behavior at a finite set of places. Therefore what really matters is the choice of the basic function $\Phi_v^0 \in \mathcal{S}(X(F_v))$.

1.3. Local conjecture of Sakellaridis–Venkatesh. In order to state the conjectural relation between X -period integrals and L -functions, we need to take a brief interlude into the local Plancherel theory of $L^2(X^\bullet(F_v))$, assuming that $X^\bullet(F_v)$ has a $G(F_v)$ -eigenmeasure.

Let \tilde{G}_X denote the spherical dual group of X . Let G_X denote the split Langlands dual group of \tilde{G}_X . The Whittaker–Plancherel theorem, which can be deduced from the usual Harish-Chandra Plancherel theorem [SV, 6.3], gives a direct integral decomposition

$$L^2((N_X(F_v), \psi) \backslash G_X(F_v)) \cong \int_{\text{Temp}(G_X)} \sigma^{\oplus m(\sigma)} d\mu_{G_X}(\sigma)$$

where $\text{Temp}(G_X)$ denotes tempered $G_X(F_v)$ -representations.

Now a paraphrase of the local conjecture is:

Conjecture 1.3.1 ([SV, Conjecture 16.5.1]). *One expects a map $\iota_* : \text{Temp}(G_X) \rightarrow \text{Irr}(G)$ ³ such that there is a unitary isomorphism*

$$(1.2) \quad L^2(X^\bullet(F_v)) \cong \int_{\text{Temp}(G_X)} \iota_*(\sigma)^{\oplus m(\sigma)} d\mu_{G_X}(\sigma).$$

Here ι_* sends tempered $G_X(F_v)$ -representations to (not necessarily tempered) unitary representations of $G(F_v)$. The Arthur packet is specified by the map $\check{G}_X \times \text{SL}_2 \rightarrow \check{G}$.

Remark 1.3.2. We stress the importance of the choice of Plancherel measure $d\mu_{G_X}$ here. For the purposes of this local conjecture, our choice is rather arbitrary – there is an entire measure class that works. But the particular choice to normalize with respect to the Tamagawa measure on $G_X(F_v)$ is of tantamount importance to the global conjecture 1.4.5 to follow.

For $\pi = \iota_*(\sigma)$ the decomposition (1.2) induces a canonical embedding⁴

$$M_\pi : \pi \otimes \bar{\pi} \rightarrow C^\infty(X^\bullet(F_v) \times X^\bullet(F_v))$$

At least formally, we can pair this with $\Phi_{1,v} \otimes \Phi_{2,v}$ for Schwartz functions $\Phi_1, \Phi_2 \in \mathcal{S}(X(F_v))$ to get a canonical Hermitian pairing⁵

$$(1.3) \quad \alpha_{\Phi_1, \Phi_2, \pi} : \pi \otimes \bar{\pi} \rightarrow C^\infty(X^\bullet(F_v) \times X^\bullet(F_v))^{\Phi_1 \otimes \Phi_2} \mathbb{C}.$$

It's not clear this pairing converges, but it's expected to almost everywhere.

Remark 1.3.3. We can write $\alpha_{\Phi_1, \Phi_2, \pi} = \ell_{\Phi_1, \pi} \otimes \ell_{\Phi_2, \bar{\pi}}$ where $\ell_{\Phi, \pi} : \pi \rightarrow \mathbb{C}$ is a linear functional

$$\ell_{\Phi, \pi} : \pi \hookrightarrow L^2(X^\bullet(F_v)) \xrightarrow{\Phi} \mathbb{C}$$

induced by (1.2), but the embedding $\pi \hookrightarrow L^2(X^\bullet(F_v))$ is not quite canonical (we can modify by an isometry). In the fortuitous circumstance where X^\bullet “unfolds” to Whittaker model (cf. [SV, Theorem 9.5.9]), there is a canonical choice of $\ell_{\Phi, \pi}$, and it becomes reasonable to compare the X -period to $\ell_{\Phi, \pi}$ without “squaring”.

1.3.4. When X is *strongly tempered* (cf. [SV, 6.2]), a condition which implies that $\check{G}_X = \check{G}$, then $M_\pi : \pi \otimes \bar{\pi} \rightarrow C^\infty(X^\bullet(F_v) \times X^\bullet(F_v))$ is defined by extension by zero from the $(G \times G)(F_v)$ -orbit of $\Delta X^\bullet(F_v) \subset X^\bullet(F_v) \times X^\bullet(F_v)$ by

$$M_\pi(u \otimes w)(x, x) = \int_{G_x} \langle \pi(h)u, w \rangle dh.$$

where G_x is the $G(F_v)$ -stabilizer of x . The fact that this integral converges and agrees with the Plancherel theorem description above is one of the main theorems [SV, Theorem 6.2.1] of Sakellaridis–Venkatesh. In particular, they show that a Plancherel decomposition like (1.2) exists when X is strongly tempered (but they don't precisely determine the multiplicities).

³To be precise, $\text{Irr}(G)$ and $L^2(X^\bullet(F_v))$ should be replaced by direct sum over all pure inner forms of X^\bullet and $\text{Temp}(G_X)$ should be replaced by the set of tempered Langlands parameters for \check{G}_X , and ι_* is a multi-valued function corresponding to Vogan Arthur packets of \check{G} .

⁴If ι_* is not injective, which means $\check{G}_X \rightarrow \check{G}$ is not injective, this embedding depends on σ not just π .

⁵Our notation α should not be confused with that of [II]; their α is what we denote α^b .

Example 1.3.5. Suppose $X = H \backslash G$ is affine homogeneous and strongly tempered. (This is the case for the Gross–Prasad variety $\mathrm{SO}_n \backslash \mathrm{SO}_n \times \mathrm{SO}_{n+1}$.) Then Φ_v can be taken to be an appropriate smoothening of the delta function $\mathbf{1}_{x_0}$ for some base point $x_0 \in X(F_v)$. Then $\alpha_{\mathbf{1}_{x_0}, \mathbf{1}_{x_0}, \pi}(u \otimes w)$ is just evaluation of $M_\pi(u \otimes w)$ at (x_0, x_0) , so we get

$$\alpha_{\mathbf{1}_{x_0}, \mathbf{1}_{x_0}, \pi}(u \otimes w) = \int_{H(F_v)} \langle \pi(h)u, w \rangle dh.$$

This is the familiar pairing that shows up in the original Ichino–Ikeda conjecture.

1.4. Generalized Ichino–Ikeda conjecture. We now explain the conjecture of Sakellaridis–Venkatesh on the relation between X -period integrals and special values of L -functions. They only state the conjecture when X is affine homogeneous, so I am taking some liberties here.

There are probably hidden assumptions needed for these conjectures to make sense: for example, X should not have type N roots⁶.

1.4.1. Let $\pi = \otimes_v \pi_v$ be a tempered (i.e., π_v is tempered at every place) and cuspidal⁷ $G(\mathbb{A})$ -representation (with an embedding into $\mathcal{A}_0(G)$).

Definition 1.4.2. For $f \in \pi$, we can formally write the X -period integral

$$(1.4) \quad \mathcal{P}_\Phi(f) := \int_{[G]} \Theta_X(\Phi, g) \cdot f(g) dg.$$

However when X is not affine homogeneous, this integral may diverge. So we *conjecturally assume* that $\mathcal{Z}(\chi, \Phi, f)$ has meromorphic continuation to $\chi = 1$ and understand $\mathcal{P}_\Phi(f)$ to mean $\mathcal{Z}(1, \Phi, f)$ in the sense of meromorphic continuation in what follows.

Ignoring convergence issues, we formally have $\mathcal{Z}(\chi, \Phi, f) = \mathcal{P}_\Phi(\chi \cdot f)$, so relating the zeta integral to an L -function is roughly equivalent to relating the X -period to a special value of an L -function. Hence in what follows we will only discuss the period and not the zeta integral, but the reader should think of the two as equivalent.

1.4.3. *Multiplicity one assumption.* For psychological purposes, from now on we will assume that for all place v , we have

$$\dim \mathrm{Hom}_{G(F_v)}(\pi_v, C^\infty(X^\bullet(F_v))) \leq 1.$$

⁸ In practice this is usually a hard theorem. Jacquet [J] also showed that even when the multiplicity one condition above fails, e.g., in the case $X = U_n \backslash \mathrm{GL}_n$, the period integral can still admit Euler product.

A consequence of the multiplicity ≤ 1 assumption is that \check{G}_X is a subgroup of \check{G} (*a priori* $\check{G}_X \rightarrow \check{G}$ is finite).

1.4.4. Let $\Phi_i = \otimes_v \Phi_{i,v} \in \mathcal{S}(X(\mathbb{A}))$ be Schwartz functions for $i = 1, 2$. We can consider the period $\mathcal{P}_\Phi|_\pi : \pi \rightarrow \mathbb{C}$ as a linear functional on π .

Assume that at each place v , the $G(F_v)$ -representation π_v occurs in the Plancherel decomposition of $L^2(X^\bullet(F_v))$. Here is where the local L^2 -theory comes in: (1.3) gives a canonical pairing $\alpha_{\Phi_1, \Phi_2, \pi_v} : \pi_v \otimes \bar{\pi}_v \rightarrow \mathbb{C}$.

⁶ \mathbb{N} is for normalizer. We want to avoid examples like $\mathrm{O}_n \backslash \mathrm{GL}_n$, which Jacquet, Mao have shown has some metaplectic behavior which is not expected to be related to L -functions.

⁷If π is a tempered representation that is not cuspidal, one can hope that there exists a regularization of the period integral in good cases.

⁸When $X \neq X^\bullet$, the space $C^\infty(X^\bullet(F_v))$ may need to be replaced by something else.

Conjecture 1.4.5 (Global conjecture, [SV, Conjecture 17.4.1]). *Let π be a tempered and cuspidal⁹ automorphic representation. For $\Phi_1, \Phi_2 \in \mathcal{S}(X(\mathbb{A}))$ we have*

$$(\mathcal{P}_{\Phi_1} \otimes \mathcal{P}_{\Phi_2})|_{\pi \otimes \bar{\pi}} = r_\phi \prod_v^* \alpha_{\Phi_1, \Phi_2, \pi_v}$$

where $r_\phi \in \mathbb{Q}^\times$ is a rational number depending on the “global Langlands parameter” ϕ associated to σ , where vaguely $\pi = \iota_*(\sigma)$.

According to [BZSV], the number r_ϕ should equal $\frac{1}{|S_\phi|}$ times the number of fixed points of the Langlands parameter acting on the dual Hamiltonian space $\check{M} = \check{G} \times^{\check{G}^X} V_X$ (see Remark 1.5.2), where S_ϕ is the centralizer in \check{G}_X of ϕ .

Remark 1.4.6. The horospherical variety $X = \overline{N \backslash G}$ fails the multiplicity one assumption, and the X -period corresponds to integrating against the constant term of an automorphic form. The constant term does *not* factorize into an Euler product, so this is an indication that at least some kind of assumption is necessary for the global conjecture to hold.

Perhaps the most reasonable general class of spherical varieties X to consider first are affine varieties that are strongly tempered and wavefront.

1.5. Understanding the Euler product. The Euler product $\prod_v^* \alpha_{\Phi_1, \Phi_2, \pi_v}$ does not converge above, so to make sense of it we need to normalize the $\alpha_{\Phi_1, \Phi_2, \pi_v}$ by certain L -factors so get the product to converge.

This is where the connection to L -functions comes in: at all but finitely many places, $\Phi_v = \Phi_v^0$ is the basic function in $\mathcal{S}(X(F_v))$ and π_v is an unramified principal series representation.

Conjecture 1.5.1. *There exists a \mathbb{Z} -graded finite-dimensional representation $\rho_X = \bigoplus_d \rho_X^d$ of the spherical L -group¹⁰ ${}^L G_X := \check{G}_X \rtimes W_{F_v}$ (where W_{F_v} is the Weil group) such that if*

- Φ_v^0 is the “IC function” associated to $X(\mathcal{O}_v)$,
- $\pi_v = \iota_*(\sigma_v)$ where σ_v (and π_v) are unramified principal series,
- $u \in \pi_v$ is $G(\mathcal{O}_v)$ -invariant vector normalized by $\|u_v\|^2 = 1$;

then one has

$$\alpha_{\Phi^0, \Phi^0, \pi_v}(u, u) = L_{X,v}^\#(1/2, \sigma_v) := \Delta_v(0) \cdot \frac{L_{X,v}(1/2, \sigma_v)}{L(1, \sigma_v, \text{Ad}_{\check{G}_X})}$$

where $\Delta_v(s)$ is a product of local zeta factors which depends only on X and not the representation σ_v , and

$$L_{X,v}(s, \sigma_v) := \prod_d L_v(s + \frac{d-1}{2}, \sigma_v, \rho_X^d).$$

Given this conjecture, we can normalize $\alpha_{\Phi_1, \Phi_2, \pi_v}$ at every place by:

$$\alpha_{\Phi_1, \Phi_2, \pi_v}^b = \frac{1}{L_{X,v}^\#(1/2, \sigma_v)} \cdot \alpha_{\Phi_1, \Phi_2, \pi_v}.$$

⁹In general, the tempered and cuspidal assumptions are both too strong in the cases of X with $P(X) \neq B$. For example, in those cases there can be no cuspidal contribution and/or there are non-tempered contributions. A less restrictive assumption is for π to have a *discrete* global Langlands parameter and an Arthur SL_2 coming from the distinguished $\check{G}_X \times \text{SL}_2 \rightarrow \check{G}$.

¹⁰There is currently no general definition of the Weil group action on \check{G}_X , so we only state the L -group for ideological completeness.

Now part of Conjecture 1.4.5 is the requirement that the global L -function $L_X^\#(s, \sigma) = \prod_v L_{X,v}^\#(s, \sigma_v)$, which is *a priori* defined for $\operatorname{Re}(s) \gg 0$, can be meromorphically continued so that one can evaluate at $s = 1/2$. Given this, Conjecture 1.4.5 says that

$$(\mathcal{P}_{\Phi_1} \otimes \mathcal{P}_{\Phi_2})|_{\pi \otimes \bar{\pi}} = r_\phi \cdot L_X^\#(1/2, \sigma) \cdot \prod_v \alpha_{\Phi_1, \Phi_2, \pi_v}^b.$$

Remark 1.5.2. The conjecture of [BZSV] says that we should further expect the representation (ρ_X, V_X) to satisfy the property that $\check{G} \times^{\check{G}^x} V_X$ is a \check{G} -Hamiltonian variety.

2. EXAMPLES

Before stating our results, we summarize some old and new examples of how this correspondence $X \rightsquigarrow \rho_X$ looks like.

2.1. Classical examples. In the table below, $T^*V = V \oplus V^*$. The names signify who discovered the corresponding integrals and determined what ρ_X should be.

	$X \circlearrowleft G$	\check{G}_X	ρ_X
Usual Langlands	Group $G' \circlearrowleft G' \times G' = G$	\check{G}'	$\check{\mathfrak{g}}'$
Whittaker normalization	$(N, \psi) \backslash G$	\check{G}	0
Tate's thesis	$\mathbb{A}^1 \circlearrowleft \mathbb{G}_m$	\mathbb{G}_m	$T^*\mathbb{C}$
Hecke	$\mathbb{G}_m \backslash \operatorname{PGL}_2$	$\check{G} = \operatorname{SL}_2$	$T^*\operatorname{std}$
Rankin–Selberg, Jacquet–Piatetski- Shapiro–Shalika	$\operatorname{GL}_n \times \mathbb{A}^n \circlearrowleft \operatorname{GL}_n \times \operatorname{GL}_n$, $H = \text{diagonal mirabolic}$	\check{G}	$T^*(\operatorname{std} \otimes \operatorname{std})$
<i>loc cit.</i>	$\operatorname{GL}_n \backslash \operatorname{GL}_{n+1} \times \operatorname{GL}_n$	\check{G}	$T^*(\operatorname{std} \otimes \operatorname{std})$
Gan–Gross–Prasad	$\operatorname{SO}_{2n} \backslash \operatorname{SO}_{2n+1} \times \operatorname{SO}_{2n}$	$\check{G} = \operatorname{SO}_{2n} \times \operatorname{Sp}_{2n}$	$\operatorname{std} \otimes \operatorname{std}$
Jacquet, Ichino	$\operatorname{PGL}_2^{\operatorname{diag}} \backslash \operatorname{PGL}_2^{\times 3}$	$\check{G} = \operatorname{SL}_2^{\times 3}$	$\operatorname{std} \otimes \operatorname{std} \otimes \operatorname{std}$

A more extensive list of examples may be found at <https://www.jonathanpwang.com/notes/RelativeDualitydb.html>

2.2. New examples. Smooth affine spherical varieties over \mathbb{C} have all been classified by [KSt]. It may still be fruitful to peruse their tables to find new examples of number theoretic interest. However, when one compares this classification with the full classification of spherical varieties, one finds that there are many more affine spherical varieties that are singular.

By considering these singular affine spherical varieties, one can find new examples number theorists haven't discovered before:

Example 2.2.1 ([SW, Example 1.1.3]). A new family of examples was provided by Sakellaridis [Sa12] generalizing the Rankin–Selberg convolution to an integral representation of the n -fold tensor product L -function for GL_2 .

Let $G = (\mathbb{G}_m \times \operatorname{SL}_2^{\times n}) / \mu_2^{\operatorname{diag}} = \operatorname{GL}_2 \times_{\det} \cdots \times_{\det} \operatorname{GL}_2$ acting on $X^\bullet = H_0 \backslash \operatorname{SL}_2^{\times n}$ where

$$H_0 = \left\{ \left(\begin{array}{cc} 1 & \\ x_1 & 1 \end{array} \right) \times \left(\begin{array}{cc} 1 & \\ x_2 & 1 \end{array} \right) \times \cdots \times \left(\begin{array}{cc} 1 & \\ x_n & 1 \end{array} \right) \mid x_1 + x_2 + \cdots + x_n = 0 \right\},$$

where $a \in \mathbb{G}_m$ acts as left multiplication by $\begin{pmatrix} a^{-1} & \\ & a \end{pmatrix}$. Let X be the affine closure of X^\bullet .

In this case $\check{G}_X = \check{G} = \operatorname{GL}_2 \times_{\det} \cdots \times_{\det} \operatorname{GL}_2$ and it follows from our work that

$$\rho_X = T^*(\operatorname{std}_2^{\otimes n} \otimes \operatorname{std}_1).$$

In fact, this is a case where the X -period \mathcal{P}_{X, Φ^0} unfolds to the Whittaker period.

Proposition 2.2.2 ([SW, Proposition 9.3.1]). *Let $\Phi^0 = \prod \Phi_v^0$ be the product of basis functions. If $f \in \pi$ is everywhere unramified and normalized so first Whittaker coefficient $W_{f,v}(1) = 1$ at all places, then*

$$\mathcal{Z}(|\det|^s, \Phi^0, f) = L(s + 1 - \frac{n}{2}, \pi, \text{std}_2^{\otimes n}).$$

for $\text{Re}(s) \gg 0$.

Our methods do not give any meromorphic continuation results about the zeta integral.

2.2.3. Explicit description of X . Let $\text{SL}_2^n = \prod^n \text{Sp}(V_i)$ where V_i are 2-dimensional symplectic spaces with fixed symplectic forms ω_i . Then the space X can be described as the set of (\underline{v}, ζ) where

- $\underline{v} = (v_i)_i \in \prod V_i$,
- $\zeta \in \wedge^n(V_1 \oplus \cdots \oplus V_n)$, such that:
- $\zeta \wedge \eta = 0$ for any $\eta = \sum c_i v_i$ with $\sum c_i = 0$,
- $\zeta \wedge v_n = \sum_i (-1)^i v_1 \wedge v_2 \wedge \cdots \wedge \omega_i \wedge \cdots \wedge v_n$, where ω_i is the fixed symplectic form on V_i . (The condition is really symmetric in the v_i 's, up to sign.)

The open X^\bullet is the subset where the v_i 's are all nonzero.

Example 2.2.4. Here is a conjectural attempt to generalize the previous Example 2.2.1. I have not checked the combinatorics, and the formulation is by observation from rank 2 cases in Wasserman's table [Was].

Let G_0 be a simply connected split semisimple group. Let N_0 denote the unipotent radical of a Borel, and define H_0 to be the kernel of a non-degenerate character $N_0 \rightarrow \mathbb{G}_a$. Define $X = \overline{H_0} \backslash G_0^{\text{aff}}$. Let $G = \mathbb{G}_m \times^{\mu_2} G_0$ act on X where \mathbb{G}_m acts on the left by $2\check{\rho}_{G_0}$, the cocharacter corresponding to the sum of positive coroots. Here $\mu_2 \rightarrow G_0$ is trivial if $\check{\rho}_{G_0}$ is in the coweight lattice. (For example when $G_0 = \text{SL}_3$ we have $G = \mathbb{G}_m \times \text{SL}_3$ where \mathbb{G}_m acts by $\check{\rho}_{\text{SL}_3}$.) Then for this (G, X) , I predict that

$$\rho_X = T^*(V^{\check{\rho}_{G_0}})$$

where $V^{\check{\rho}_{G_0}}$ really means the highest weight representation of \check{G} with highest weight corresponding to $\mathbb{G}_m \rightarrow G : a \pmod{\mu_2} \mapsto (a, (2\check{\rho}_{G_0})(a)) \pmod{\mu_2}$.

3. SUMMARY OF RESULTS

A rough summary of the content of [SW] is that we “almost” prove Conjecture 1.5.1 over a local function field in the case where $\check{G}_X = \check{G}$ and X has no spherical roots of type N.

3.1. Previous work. When $X = H \backslash G$ is *affine homogeneous* (by [Lun73], [Ric77], this is equivalent to H being reductive), Sakellaridis ([Sa08, Sa13]) proved¹¹ Conjecture 1.5.1 using function-theoretic techniques. In these cases, X is smooth so Φ_0 is just the indicator function of $X(\mathcal{O}_v)$ in $X(F_v)$.

However when X is singular, geometric considerations are needed if we want to understand the relation between intersection complexes (perverse sheaves) and L -values.

Explicit formulas for the “IC function” have been previously established (as well as other geometric results) in the following cases:

- Braverman–Finkelberg–Gaitsgory–Mirković [BFGM]:
 - $X = \overline{N} \backslash \check{G}$, $\check{G}_X = \check{T}$, $V_X = \check{\mathfrak{g}}^*/\check{\mathfrak{t}}^*$. Note that $\check{G} \times^{\check{G}_X} V_X = T^*(\check{G}/\check{T})$, which has some global incarnation as geometric Eisenstein series

¹¹Technically, his work gives ρ_X as a *virtual* representation but didn't show it's a true representation.

- Bouthier–Ngô–Sakellaridis [BNS]:
 - $X \supset G'$ is an L -monoid, so here the group is $G = G' \times G'$, $\check{G}_X = \check{G}'$, and $V_X = \check{\mathfrak{g}}' \oplus T^*V^\lambda$ where λ is the coweight appearing in the definition of an L -monoid.

3.2. Main result on Plancherel measure. From now on, we will work entirely in the local setting and drop the subscript v . Let $F = \mathbb{F}_q((t))$ denote a local function field and $\mathcal{O} = \mathbb{F}_q[[t]]$ its ring of integers. We allow G to be a (not necessarily split) connected reductive group over \mathbb{F}_q . Let X be an affine spherical variety over \mathbb{F}_q .

3.2.1. In the unramified situation, the map ι_* is more transparent: an unramified $G_X(F)$ -representation is a principal series, the induction of an unramified character on $T_X(F)$. Such a character corresponds to a point in $\check{T}_X(\mathbb{C})$. Likewise, an unramified $G(F)$ -representation π corresponds to a point in $\check{T}(\mathbb{C})$. The map ι_* on unramified representations is just the map of dual maximal tori $\check{T}_X \rightarrow \check{T}$ induced by $\check{G}_X \rightarrow \check{G}$.

3.2.2. Suppose $\check{G}_X = \check{G}$ and π is an unramified $G(F)$ -representation. Then π is the induction of a character $\chi : T(F) \rightarrow \mathbb{C}$. Note that the local L -factor $L(s, \chi, \rho'_X)$ can be defined for any \check{T} -representation ρ'_X , without requiring ρ'_X to be a \check{G} -representation. With this, we can state our main theorem:

Theorem 3.2.3 (Sakellaridis–W). *Assume X affine spherical, $\check{G}_X = \check{G}$ and X has no type N roots¹². Then there is a $(\check{T} \rtimes \langle \text{Fr} \rangle)$ -representation ρ'_X such that: for $u \in \pi_\chi$ a $G(\mathcal{O})$ -invariant vector normalized by $\|u\|^2 = 1$, we have*

$$\alpha_{\Phi^0, \Phi^0, \pi_\chi}(u, u) = \Delta(0) \frac{L(1/2, \chi, \rho'_X)}{L(1, \chi, \text{Ad})}$$

The \check{T} -representation ρ'_X has the properties:

- (i) ρ'_X has an action of $(\text{SL}_2)_\alpha$ for every simple root α
 - We do not check the Weyl/Serre relations, which would imply ρ'_X is a \check{G} -representation.
- (ii) Assuming ρ'_X is a \check{G} -representation, we determine its highest weights with multiplicities (in terms of prime B -divisors of X).

The factor $\Delta(0)$ can be extracted from [SV, (17.8)].

We are confident that ρ'_X is indeed a \check{G} -representation, but for clarity let us call ρ_X the \check{G} -representation that it would have to equal according to (ii) above. Below we explain the definition of the assignment $X \rightsquigarrow \rho_X$.

Remark 3.2.4. Note that for a given X , if ρ_X is minuscule, then there is no question of weight multiplicities, so ρ'_X must equal ρ_X , i.e., we prove Conjecture 1.5.1 in those cases.

We also show that if X is affine homogeneous, then ρ_X must be minuscule.

We can also reduce the checking of $\rho'_X = \rho_X$ to the cases where $X = \overline{H \backslash G}^{\text{aff}}$ and G has semisimple rank 2. There are only 5-6 such cases with ρ_X non-minuscule (four of which come from Example 2.2.4 for types A_2, B_2, C_2, G_2), which can in principle be checked “by hand”.

There is some hope that our techniques will generalize to any X (no restriction on \check{G}_X) by combining the knowledge from [BFGM, BNS].

¹²We also need some further assumptions over \mathbb{F}_q to ensure X behaves like it does in characteristic 0

3.3. Recipe for ρ_X . We now explain the combinatorial recipe for how to determine ρ_X from X . The pattern is expected to hold for any X without type N roots but we only prove it so far when $\check{G}_X = \check{G}$. We base change X from \mathbb{F}_q to $k = \overline{\mathbb{F}}_q$ below.

The combinatorial data attached to X has two parts:

- (a) one comes from the open G -orbit X^\bullet and
- (b) the other comes from the affine embedding $X^\bullet \hookrightarrow X$.

Part (b) gives a finite collection of anti-dominant weights $\check{\theta}_i \in \Lambda_{\check{G}_X}^-$. Then

$$\rho_X = \rho_{X^\bullet} \oplus \bigoplus T^* V^{\check{\theta}_i}$$

where $V^{\check{\theta}_i}$ is the irreducible \check{G}_X -representation with lowest weight $\check{\theta}_i$ and ρ_{X^\bullet} is a representation determined by the data from (a) that we now describe.

In part (a), we consider the prime B -stable divisors in X^\bullet . These are called *colors*. A color D determines a valuation v_D on $k(X^\bullet)$ and in particular on the B -eigenvectors $k(X^\bullet)^{(B)}$. Since X is spherical, $k(X^\bullet)^{(B)}/k^\times = \check{\Lambda}_{\check{G}_X}$, the coweight lattice of \check{T}_X . Thus restricting v_D to $k(X^\bullet)^{(B)}$ determines a weight $\check{\nu}_D \in \Lambda_{\check{G}_X}$. Now ρ_{X^\bullet} is the unique finite dimensional \check{G}_X -representation such that

- The highest weights of ρ_{X^\bullet} are precisely the set $\Lambda_{\check{G}_X}^+ \cap W_X \{\check{\nu}_D\}_{\text{colors } D}$, where W_X is Weyl group of \check{G}_X , with multiplicities determined by:
- The multiplicity of the weight space of weight $\check{\nu}_D$ in ρ_{X^\bullet} is equal to the number of colors D' such that $\check{\nu}_D = \check{\nu}_{D'}$ (this number is either 1 or 2).

The graded degree of ρ_{X^\bullet} is 1 (so corresponds to central L -value at $1/2$) while the degree of $T^* V^{\check{\theta}_i}$ is a constant determined by $\check{\theta}_i$ and the G -eigen-volume form on $X^\bullet(F)$.

Remark 3.3.1. The case when two colors D, D' have $\check{\nu}_D = \check{\nu}_{D'}$ arises from situations where you have $\mathbb{G}_m \backslash \text{PGL}_2$ instead of $\mathbb{G}_m \backslash \text{GL}_2$. In this case $\check{\nu}_D = \check{\alpha}/2$ for a simple root $\check{\alpha}$ of \check{G}_X . In practice you can always replace G by a central extension to remove this case.

3.4. Assumption $\check{G}_X = \check{G}$. The condition that X has $\check{G}_X = \check{G}$ and no type N roots has an easily accessible description, which I now explain.

This is equivalent to the following (after base change to $\overline{\mathbb{F}}_q$):

- X has an open B -orbit X° acted on simply transitively by B (so after choosing a base point $x_0 \in X^\circ$ we get $X^\circ \cong B$),
- $X^\circ P_\alpha / \mathcal{R}(P_\alpha) \cong \mathbb{G}_m \backslash \text{PGL}_2$ for every simple α . Here $P_\alpha \supset B$ is the standard sub-minimal parabolic corresponding to α .

So this says X has open subvarieties which “look” like the Hecke case $\mathbb{G}_m \backslash \text{GL}_2$, and the complement of these opens are certain B -divisors.

Remark 3.4.1. An important corollary of this assumption is that the generic stabilizer subgroup H is *connected*. This assumption saves us from many headaches/complications at various places. Geometrically, one would need to consider moduli of $\pi_0(H)$ -local systems at various places.

3.5. Bernstein asymptotics. The computation of $\alpha_{\Phi^0, \Phi^0, \pi_X}(u, u)$ is done using the theory of Bernstein asymptotics developed in [SV]. We give a very brief idea of how this goes, without any assumption on \check{G}_X .

3.5.1. Plancherel decomposition (1.2) gives a decomposition

$$\|\Phi^0\|^2 = \int_{\text{Temp}(G_X)} \|\Phi^0\|_\sigma^2 d\mu_{G_X}(\sigma).$$

Since Φ^0 is $G(\mathcal{O})$ -invariant, we will only get contributions on the right hand side from unramified σ . In this formulation we have

$$\alpha_{\Phi^0, \Phi^0, \pi}(u, u) = \|\Phi^0\|_{\sigma}^2,$$

where $\iota_*(\sigma) = \pi$ is unramified.

3.5.2. From the usual Harish-Chandra theory of “parabolic descent”, we have a direct sum decomposition

$$L^2((N_X(F), \psi) \backslash G_X(F)) = \bigoplus_{\Theta} L^2((N_X(F), \psi) \backslash G_X(F))_{\Theta, \text{disc}}$$

ranging over conjugacy classes of parabolic subgroups of G_X , or equivalently subsets of Δ_X , the set of simple coroots of \check{G}_X (= the set of spherical roots of X).

Sakellaridis–Venkatesh establish the parallel decomposition for $L^2(X^{\bullet}(F))$.

Theorem 3.5.3 ([SV, Scattering Theorem 7.3.1]). *Suppose X^{\bullet} is wavefront and satisfies some technical assumptions. Then there is a decomposition*

$$L^2(X^{\bullet}(F)) = \bigoplus_{\Theta} L^2(X^{\bullet}(F))_{\Theta}$$

running over subsets $\Theta \subset \Delta_X$, where roughly speaking $L^2(X^{\bullet}(F))_{\Theta}$ is the image of a canonical G -equivariant “ L^2 Bernstein morphism”

$$\iota_{\Theta} : L^2(X_{\Theta}^{\bullet}(F)) \rightarrow L^2(X^{\bullet}(F))$$

and X_{Θ} is a boundary degeneration of X at “ Θ -infinity”.

Here ι_{Θ} can be thought of as an analog of unitary Eisenstein series (or Harish-Chandra’s Eisenstein integral).

Luckily for us, we can show $\|\Phi^0\|^2$ only has contributions from the most continuous spectrum, which corresponds to $\Theta = \emptyset$. Thus we have

$$\|\Phi^0\|^2 = \int_{\check{T}_X^1/W_X} \|\Phi^0\|_{\chi}^2 d\chi.$$

where \check{T}_X^1 denotes unitary unramified characters of $T_X(F)$. Here $d\chi$ is slightly different from $d\mu_{G_X}(\sigma_{\chi})$, which will contribute a normalization factor of $\frac{L(1, \chi, \text{Ad})}{L(0, \chi, \text{Ad})}$. From Theorem 3.5.3 it follows that $\|\Phi^0\|_{\chi}^2 = \frac{1}{|W_X|} \|\iota_{\emptyset}^* \Phi^0\|_{\chi}^2$, where $\iota_{\emptyset}^* : L^2(X^{\bullet}(F)) \rightarrow L^2(X_{\emptyset}^{\bullet}(F))$ is the adjoint of ι_{\emptyset} .

3.5.4. The map ι_{\emptyset}^* in fact comes from a “smooth Bernstein asymptotics” map

$$e_{\emptyset}^* : C^{\infty}(X^{\bullet}(F)) \rightarrow C^{\infty}(X_{\emptyset}^{\bullet}(F))$$

which is roughly characterized by being $G(F)$ -equivariant and capturing the asymptotic behavior of the function on $X^{\bullet}(F)$. Then ι_{\emptyset}^* is defined from e_{\emptyset}^* by throwing away some “exponents” that do not belong to the L^2 -space.

A global analog of e_{\emptyset}^* is the constant term functor, while the analog of ι_{\emptyset}^* is Harish-Chandra’s (L^2 -)constant term integral. See also the first paragraph of [SV, §8].

3.6. Main result in terms of asymptotics. Our result is more precisely stated in terms of asymptotics. We return to assuming $\check{G}_X = \check{G}$. In this setting, $X_\emptyset^\bullet = N^- \backslash G$.

Theorem 3.6.1 (Sakellaridis-W). *Same assumptions as in Theorem 3.2.3.*

- (i) *The asymptotics map e_\emptyset^* corresponds, under the functions-sheaves dictionary, to the nearby cycles functor on algebro-geometric models of $X^\bullet(F), X_\emptyset^\bullet(F)$.*
- (ii) *There is a \check{T} -polarization of $\rho'_X = T^*(V_X^+)$ where V_X^+ is a \mathbb{Z} -graded $\check{T} \rtimes \langle \text{Fr} \rangle$ -representation such that*

$$(3.1) \quad (\eta\delta)^{\frac{1}{2}}(t^{\check{\lambda}}) \cdot e_\emptyset^* \Phi^0(t^{\check{\lambda}}) = \frac{L(\frac{1}{2}, \chi, V_X^+)}{L(0, \chi, \check{\mathfrak{n}})}$$

Here η is the eigencharacter of the eigenmeasure on $X^\bullet(F)$. The factor $(\eta\delta)^{\frac{1}{2}}$ is a normalization factor, cf. Remark 4.1.6. The $\frac{1}{2}$ special L -value should be shifted according to the \mathbb{Z} -grading on V_X^+ , which is 0 for $V_{X^\bullet}^+$, so I suppress the notation.

3.7. Main result in terms of Radon transform. In practice, the definitions of both e_\emptyset^* and nearby cycles are rather opaque and hard to compute with. We instead do all our computations by passing to the Radon transform.

From now on we will fix a base point $x_0 \in X^\circ(\mathbb{F}_q)$ in the open B -orbit (such x_0 exists by [Sa08, Proposition 3.2.1]).

Definition 3.7.1. The X -Radon transform

$$\pi_! : C_c^\infty(X^\bullet(F)) \rightarrow C^\infty((N \backslash G)(F))$$

is defined by integrating over generic horocycles

$$\pi_! \Phi(g) := \int_{N(F)} \Phi(x_0 n g) dn, \quad g \in G(F).$$

The relation between e_\emptyset^* and $\pi_!$ is that there is a commutative diagram

$$(3.2) \quad \begin{array}{ccc} C_c^\infty(X^\bullet(F)) & \xrightarrow{e_\emptyset^*} & C^\infty(X_\emptyset^\bullet(F)) \\ & \searrow \pi_! & \swarrow \pi_{\emptyset!} \\ & C^\infty((N \backslash G)(F)) & \end{array}$$

where $\pi_{\emptyset!} : C_+^\infty((N^- \backslash G)(F)) \rightarrow C_-^\infty((N \backslash G)(F))$ is the standard long intertwining operator, which is *invertible* if one considers some spaces with suitable support conditions (cf. [BK, Proposition 7.5(b)], [W, Proposition 2.8.5]). Therefore assuming some convergence properties, we can think of e_\emptyset^* and $\pi_!$ as “the same”.

Our main result in terms of the Radon transform, from which we recover Theorem 3.6.1 is:

Theorem 3.7.2 (Sakellaridis-W). *Same assumptions and notation as in Theorem 3.6.1.*

- (i) *We show a triangle analogous to (3.2) commutes in the geometric setting of nearby cycles using the contraction principle (i.e., Braden’s theorem).*
- (ii) *The Mellin transform of $\pi_! \Phi^0 \in C^\infty((N \backslash G)(F))$ equals*

$$(3.3) \quad \widehat{\pi_! \Phi^0}(\chi) := \int_{T(F)} (\pi_! \Phi^0)(t) \chi(t) dt = \frac{L(\frac{1}{2}, \chi, V_X^+)}{L(1, \chi, \check{\mathfrak{n}})}$$

on unramified characters $\chi : T(F) \rightarrow \mathbb{C}$ where the Mellin transform is convergent.

3.7.3. Here is a more explicit description of $\pi_! \Phi^0$.

Note that $\pi_! \Phi^0 \in C^\infty((N \backslash G)(F))^{G(\mathcal{O})} = \text{Fn}(\check{\Lambda}_G)$ is a function on the set of coweights of G . Assume that all the prime B -divisors of X_k are defined over \mathbb{F}_q (I think this implies that the Frobenius action on \check{G}_X is trivial, i.e., ${}^L G_X = \check{G} \times \langle \text{Fr} \rangle$). Then (3.3) is equivalent to saying

$$(3.4) \quad \pi_! \Phi^0 = \frac{\prod_{\check{\alpha} \in \check{\Phi}_G^+} (1 - q^{-1} e^{\check{\alpha}})}{\prod_{\check{\lambda} \in \text{wt}(V_X^+)} (1 - q^{-\frac{1}{2}} e^{\check{\lambda}})}$$

where $e^{\check{\lambda}}$ is the indicator function of $\check{\lambda}$ and $e^{\check{\lambda}} e^{\check{\mu}} = e^{\check{\lambda} + \check{\mu}}$. The Euler product on the right should be understood via a power series expansion:

$$\frac{1}{1 - q^{-\frac{1}{2}} e^{\check{\lambda}}} = \sum_{n \geq 0} (q^{-\frac{1}{2}} e^{\check{\lambda}})^n.$$

3.8. **Formula for Φ^0 .** The basic function Φ^0 is a $G(\mathcal{O}_v)$ -invariant function on $X^\bullet(F)$.

We have the following parametrization of the set of orbits $X^\bullet(F)/G(\mathcal{O})$, proved by [GN] for $\mathbb{C}((t))$ and extended to non-archimedean local fields F by [Sa12].

Theorem 3.8.1 ([Sa12, Theorem 2.3.8]). *Assume X^\bullet has good integral model over \mathcal{O} . If Λ_G/Λ_X is torsion-free then there is a bijection between the set $X^\bullet(F)/G(\mathcal{O})$ and $-\Lambda_{G_X}^+ \subset \check{\Lambda}_X$.*

Our assumptions (3.4) obvious satisfy the above. Hence in our case Φ^0 is just a function on the monoid $-\check{\Lambda}_G^+$. We can in fact give a formula for Φ^0 based on a variant of ‘‘inverse Satake transform’’ for asymptotics ([Sa18, Corollary 5.5]) once we know what the representation $\rho'_X = T^* V_X^+$ is.

Proposition 3.8.2. *We consider Φ^0 as a function on $-\check{\Lambda}_G^+$ and $e_\emptyset^* \Phi^0$ as a function on $\check{\Lambda}_G$. Then*

$$(3.5) \quad \Phi^0 = (e_\emptyset^* \Phi^0)|_{-\check{\Lambda}_G^+}.$$

Since we have a formula for $e_\emptyset^* \Phi^0$ in terms of V_X^+ (see for example (3.4)), we could just use (3.5) as the definition of Φ^0 without talking about IC functions. Then you can probably establish the formula for $e_\emptyset^* \Phi^0$ *a posteriori* by classical methods. Our proof that Φ^0 equals the IC function of $X(\mathcal{O})$ is focused on establishing the connection between geometry and number theory in the hopes that this will lead to new methods in studying L -functions.

4. GEOMETRIC TECHNIQUES

Let $F = \mathbb{F}_q((t))$ and $\mathcal{O} = \mathbb{F}_q[[t]]$ as above. Let $k = \overline{\mathbb{F}_q}$; for the geometric results below, we can also take $k = \mathbb{C}$. Let $\mathbf{F} = k((t))$ and $\mathbf{O} = k[[t]]$.

4.1. **IC function.** We would like to say that the IC function Φ^0 is the trace of geometric Frobenius acting on the IC complex of $X(\mathcal{O})$.

To define IC complexes you need a theory of perverse sheaves (perverse t-structure) on an algebraic variety. Fortunately, we can give $X(\mathcal{O})$ algebro-geometric structure as follows:

Definition 4.1.1. The *formal arc space* of X is the scheme $\mathbf{X}_{\mathbf{O}}$ over \mathbb{F}_q with functor of points

$$\mathbf{X}_{\mathbf{O}}(R) = X(R[[t]])$$

for a test \mathbb{F}_q -algebra R .

By definition, the \mathbb{F}_q -points of $\mathbf{X}_{\mathbf{O}}$ equal $X(\mathcal{O})$.

Remark 4.1.2. For any scheme X of finite type, $\mathbf{X}_{\mathbf{O}}$ is representable by a scheme. When X is affine, so is $\mathbf{X}_{\mathbf{O}}$. However the functor $\mathbf{X}_{\mathbf{F}}(R) = X(R((t)))$ for the *formal loop space* is only representable by an ind-scheme when X is affine – it does not have good properties for non-affine X .

4.1.3. Now we run into a problem. The theory of perverse sheaves is developed in [BBDG] for schemes (locally) of *finite type*. On the other hand, the scheme $\mathbf{X}_{\mathbf{O}}$ is very much of infinite type. So far there is no general way to define perverse sheaves on infinite type schemes.

In the situation of the affine Grassmannian $\mathrm{Gr}_G = \mathbf{G}_{\mathbf{F}}/\mathbf{G}_{\mathbf{O}}$, the $\mathbf{G}_{\mathbf{O}}$ -orbit closures are at least of finite type, which saves the day. However for general affine X , the stack quotient $\mathbf{X}_{\mathbf{O}}/\mathbf{G}_{\mathbf{O}}$ will still not have strata that look like finite type schemes.

Nevertheless, Bouthier–Ngô–Sakellaridis [BNS] show that the IC function of $\mathbf{X}_{\mathbf{O}}$, which should equal the trace of geometric Frobenius of $\mathrm{IC}_{\mathbf{X}_{\mathbf{O}}}$, is well-defined. They use a theorem of Grinberg–Kazhdan (characteristic 0) and Drinfeld (any characteristic):

Theorem 4.1.4 (Grinberg–Kazhdan, Drinfeld). *Let $\gamma \in X(\mathbb{F}_q[[t]])$ be an arc that generically lands in the smooth locus of X . Then there exists a finite type scheme Y and $y \in Y(\mathbb{F}_q)$ such that there is an isomorphism of formal neighborhoods*

$$(\widehat{\mathbf{X}_{\mathbf{O}}})_{\gamma} \cong \widehat{Y}_y \times \widehat{\mathbb{A}^{\infty}}.$$

I.e., near generic arcs $\mathbf{X}_{\mathbf{O}}$ has finite-type singularities.

We call Y as above a *model* of $\mathbf{X}_{\mathbf{O}}$.

Definition 4.1.5. The IC function Φ^0 of $X(\mathcal{O})$ is defined by

$$\Phi^0(\gamma) := \mathrm{tr}(\mathrm{Fr}_y, \mathrm{IC}_Y|_y^*)$$

where (Y, y) are as in Theorem 4.1.4. [BNS] show that this definition is independent of the choice of Y .

Remark 4.1.6. Here IC_Y is normalized *without the Tate twist*, so if Y is smooth then $\mathrm{IC}_Y = \overline{\mathbb{Q}}_{\ell Y}$. It is hard to make sense of the Tate twist since the dimension becomes infinite. However this discrepancy of normalizations is what accounts for the factor $(\eta\delta)^{\frac{1}{2}}$ in (3.1).

4.2. **Models for the formal arc space.** We will use the fact that Drinfeld’s proof [D] of this theorem gives us explicit models for $\mathbf{X}_{\mathbf{O}}$. This phenomenon was first used by Finkelberg–Mirković to study $X = \overline{G/N}$ ($\check{G}_X = \check{T}$). The two models are:

- the Artin stack $\mathcal{M}_X = \mathrm{Maps}_{\mathrm{gen}}(C, X/G \supset X^{\bullet}/G)$, which we call the *global model*¹³.
- the *Zastava space*¹⁴ $\mathcal{Y}_X = \mathrm{Maps}_{\mathrm{gen}}(C, X/B \supset X^{\circ}/B)$.

Both models are important for different reasons:

- The global model \mathcal{M}_X allows us to model the part of the Hecke action of $\mathbf{G}_{\mathbf{F}}$ on $\mathbf{X}_{\mathbf{F}}$ that stays in $\mathbf{X}_{\mathbf{O}}$. (More generally, one can define an ind-stack $\mathcal{M}_X^{H\text{-gen}}$ of maps with poles that models the $\mathbf{G}_{\mathbf{F}}$ -action on $\mathbf{X}_{\mathbf{F}} - (\mathbf{X} - \mathbf{X}^{\bullet})_{\mathbf{F}}$.)
- The Zastava space \mathcal{Y}_X has a graded *factorization property* that is key to making the connection with L -values.

Drinfeld’s proof [D] of Theorem 4.1.4 directly shows that \mathcal{Y}_X is a model for $\mathbf{X}_{\mathbf{O}}$. There is a general yoga that passes from the Zastava model \mathcal{Y}_X to the global model \mathcal{M}_X ; or one can argue directly as follows:

¹³In certain places in the literature this is called the space of quasi-maps

¹⁴Zastava is Croatian for flag

Fix a point $v \in C(\mathbb{F}_q)$. Identify the completed local ring \mathcal{O}_v at C with \mathcal{O} . Now we have a map $\text{Spec } \mathcal{O}_v \rightarrow C$. Restricting along this map gives a map

$$(4.1) \quad \mathcal{M}_X = \text{Maps}_{\text{gen}}(C, X/G \supset X^\bullet/G) \rightarrow \text{Maps}(\text{Spec } \mathcal{O}_v, X/G) \cong \mathbf{X}_{\mathbf{O}}/\mathbf{G}_{\mathbf{O}}.$$

Now one can show that if you restrict (4.1) to functions $f : C \rightarrow X/G$ such that $f(C - v) \subset X^\bullet/G$, then (4.1) is formally smooth.

4.2.1. Considerations of the partial Hecke action on \mathcal{M}_X allow us to reduce the study of X to that of just $\overline{X^\bullet}^{\text{aff}}$. (This is a simplification of some technical results, whose proofs require inspecting the interactions between the global and Zastava models.)

4.2.2. The global model is also of interest because studying the pushforward of $\text{IC}_{\mathcal{M}_X}$ along the map $\mathcal{M}_X \rightarrow \text{Bun}_G$ using geometric methods is expected to give new approaches to the relative trace formula associated to X (cf. [N, §3]).

4.3. Zastava space and the graded factorization property. From now on I will base change to k while keeping the same notation. (Everything is canonical enough that the action of Frobenius is easy to keep track of.)

We fix a base point $x_0 \in X^\circ(k)$ and identify $X^\circ \cong B$ under assumption 3.4. Our assumptions imply that the stack X/B contains $X^\circ/B = \text{pt}$ as an open substack.

4.3.1. A point $y \in \mathcal{Y}_X(k)$ is a map $C \rightarrow X/B$ generically landing in pt . So by Beauville–Laszlo’s theorem

$$y \leftrightarrow \left\{ \begin{array}{l} \text{finite set } \{v_i\}_{i \in I} \subset C(k), \\ \hat{y}_i \in (X(\mathcal{O}_{v_i}) \cap X^\circ(F_{v_i}))/B(\mathcal{O}_{v_i}), \\ y(C - \{v_i\}) = \text{pt} \end{array} \right\}$$

Recall we are using $x_0 \in X^\circ(k)$ to identify $X^\circ \cong B$. Then

$$X^\circ(F_{v_i})/B(\mathcal{O}_{v_i}) \cong \mathbf{B}_{\mathbf{F}_{v_i}}/\mathbf{B}_{\mathbf{O}_{v_i}}(k) = \text{Gr}_{B, v_i}(k)$$

Now recall that Gr_B has the same connected components as Gr_T , which are indexed by the coweight lattice $\check{\Lambda}$. So to each \hat{y}_i is attached a coweight $\check{\lambda}_i \in \check{\Lambda}$.

From this we see that \mathcal{Y} lives over a *configuration space*

$$\left\{ \check{\Lambda}\text{-valued divisors : } \sum_{i \in I} \check{\lambda}_i \cdot v_i, v_i \in C(k) \text{ distinct} \right\}$$

If $\check{\lambda}_i$ could be any coweight then we would need something fancy like the Ran space to make sense of the above. However, since $\hat{y}_i \in X(\mathcal{O}_{v_i})$ is an arc, all the $\check{\lambda}_i$ belong to a strictly convex cone. So there is a sense of “positive” grading. More specifically,

$$\pi : \mathcal{Y} \rightarrow \mathcal{A} = \text{Maps}(C, X//N/T).$$

Let me assume for ultimate simplicity that $X//N = \mathbb{A}^r \supset \mathbb{G}_m^r = T$ with a corresponding basis $\check{\nu}_1, \dots, \check{\nu}_r \in \check{\Lambda}$ for the cocharacters whose limit as $t \rightarrow 0$ lands in $X//N$. Then

$$\mathcal{A} = \text{Maps}(C, \mathbb{A}^r/\mathbb{G}_m^r) = (\text{Sym } C)^r = \bigsqcup_{(n_i) \in \mathbb{N}^r} C^{(n_1)} \times \dots \times C^{(n_r)} =: \bigsqcup \mathcal{A}^{n_1 \check{\nu}_1 + \dots + n_r \check{\nu}_r}$$

is the scheme of r divisors on C . Let the preimage of $\mathcal{A}^{\check{\lambda}}$ be $\mathcal{Y}^{\check{\lambda}}$.

Then $\mathcal{Y}^{\check{\lambda}}$ is a *finite type* scheme.

4.3.2. *Graded factorization.* Notice that the fiber over $\check{\lambda}_1 \cdot v_1 + \check{\lambda}_2 \cdot v_2 \in \mathcal{A}^{\check{\lambda}_1 + \check{\lambda}_2}$ where v_1, v_2 are distinct, only depends on the independent fibers over $\check{\lambda}_1 \cdot v_1$ and $\check{\lambda}_2 \cdot v_2$. This is called a *graded factorization* property of (the collection of components of) \mathcal{Y} .

Aside: in the situation above the $\mathcal{Y}^{\check{\lambda}}$ are indeed irreducible components, but we could only prove this in a *very* roundabout way.

4.4. **Central fibers.** The graded factorization property essentially says the fiber of π over $\check{\lambda} \cdot v$ at a single point $v \in C(k)$ is the most important. This fiber is isomorphic to

$$\mathcal{Y}^{\check{\lambda}} := \mathrm{Gr}_{B,v}^{\check{\lambda}} \times_{\mathbf{X}_{\mathbf{F}}/\mathbf{B}_{\mathbf{O}}} \mathbf{X}_{\mathbf{O}}/\mathbf{B}_{\mathbf{O}},$$

where $\mathbf{B}_{\mathbf{F}} \rightarrow \mathbf{X}_{\mathbf{F}}$ is the action on x_0 . This fiber doesn't depend on v . Observe that

$$\mathrm{tr}(\mathrm{Fr}, \pi_! \mathrm{IC}_{\mathcal{Y}}|_{\check{\lambda} \cdot v}^*) = \mathrm{tr}(\mathrm{Fr}, H_c^*(\mathcal{Y}^{\check{\lambda}}, \mathrm{IC}_{\mathcal{Y}})) = \int_{N(F)} \Phi_0(x_0 n t^{\check{\lambda}}) = \pi_! \Phi_0(t^{\check{\lambda}})$$

is the Radon transform we wanted to calculate back in (3.4).

Example 4.4.1. Let $X = \mathbb{G}_m \backslash \mathrm{GL}_2$ where $\mathbb{G}_m = (*_1)$. Then $\mathcal{Y} = \mathrm{Maps}_{\mathrm{gen}}(C, X/B) = \mathrm{Maps}_{\mathrm{gen}}(C, \mathbb{G}_m \backslash \mathbb{P}^1)$ parametrizes

$$\mathcal{A}, \mathcal{L} \in \mathrm{Pic}, \mathcal{L} \xrightarrow{(x,y)} \mathcal{A} \oplus \mathcal{O}.$$

Generically landing in X° means x, y do not simultaneously vanish after taking fiber at any point. What this amounts to is two divisors with disjoint support:

$$\mathcal{Y} = \mathrm{Sym} C \overset{\circ}{\times} \mathrm{Sym} C$$

Meanwhile $X//N = \mathbb{A}^2$ with basis $\check{\varepsilon}_1 = (1, 0)$, $-\check{\varepsilon}_2 = (0, -1)$. So

$$\pi : \mathcal{Y} = \mathrm{Sym} C \overset{\circ}{\times} \mathrm{Sym} C \rightarrow \mathrm{Sym} C \times \mathrm{Sym} C = \mathcal{A}$$

is an open embedding. The preimage of $(n_1 \check{\varepsilon}_1 - n_2 \check{\varepsilon}_2) \cdot v$ is empty if n_1, n_2 are both nonzero, and a point otherwise. So we see

$$\pi_! \Phi_0 = e^0 + \sum_{n \geq 1} (q^{-n/2} e^{n \check{\varepsilon}_1} + q^{-n/2} e^{-n \check{\varepsilon}_2}) = \frac{1 - q^{-1} e^{\check{\alpha}}}{(1 - q^{-1/2} e^{\check{\varepsilon}_1})(1 - q^{-1/2} e^{-\check{\varepsilon}_2})}$$

since $\check{\alpha} = \check{\varepsilon}_1 - \check{\varepsilon}_2$. Note that $|\widehat{\pi_! \Phi_0}(\chi)|^2 = \frac{L(\chi, \mathrm{std} \oplus \mathrm{std}^*, 1/2)}{L(\chi, \mathfrak{g}/\mathfrak{k}, 1)}$.

As we see above, π is not proper, but we can compactify it to:

$$\overline{\mathcal{Y}} = \mathrm{Maps}(C, X \times \overline{G/N} / (G^{\mathrm{diag}} \times T))$$

and we still have $\overline{\pi} : \overline{\mathcal{Y}} \rightarrow \mathcal{A}$. Let $\overline{\mathcal{Y}}^{\check{\lambda}}$ be preimage of $\mathcal{A}^{\check{\lambda}}$; then $\overline{\mathcal{Y}}$ still has the graded factorization property.

Theorem 4.4.2 (Sakellaridis–W). *Under our assumptions on X , the map $\overline{\pi} : \overline{\mathcal{Y}} \rightarrow \mathcal{A}$ is stratified semi-small.*

We emphasize that this is extremely special to the $\check{G}_X = \check{G}$ case! The statement is definitely false for example when $X = \overline{N} \backslash \overline{G}$.

Toy situation: if $\overline{\mathcal{Y}}$ were smooth, then semi-smallness for $\overline{\pi}$ amounts to (because of factorization):

$$(4.2) \quad \dim \overline{\mathcal{Y}}^{\check{\lambda}} \leq \mathrm{crit}(\check{\lambda})$$

The general situation is more complicated because of $\mathbf{G}_{\mathbf{O}}$ -orbit strata, but using our results on the Hecke action on \mathcal{M}_X , we get roughly the same requirement on the central fibers of X^\bullet .

Under some assumptions, $\text{crit}(\check{\lambda}) = \frac{\text{len}(\check{\lambda})-1}{2}$, where $\text{len}(\check{\lambda})$ is the number of $\check{\nu}_D$, for non-distinct colors D , that sum to equal $\check{\lambda}$.

4.4.3. A fact that is presumably known to experts but not often stated is that in the above situation where you have a semi-small map, the decomposition theorem together with the graded factorization property immediately tell you that

$$(4.3) \quad \text{tr}(\text{Fr}, (\bar{\pi}_! \text{IC}_{\bar{y}})|_{?.v}^*) = \frac{1}{\prod_{\check{\lambda} \in \mathfrak{B}^+} (1 - q^{-\frac{1}{2}} e^{\check{\lambda}})}$$

has the desired Euler product format. Here \mathfrak{B}^+ corresponds to the *relevant strata* supported at a single point. More specifically, $\mathfrak{B}^+ =$ the irreducible components of $\bar{Y}^{\check{\lambda}}$ of $\dim = \text{crit}(\check{\lambda})$ as $\check{\lambda}$ varies. (This is an oversimplification but it's almost true.)

Remark 4.4.4. Note that the right hand side of (4.3) almost looks like (3.4). The difference between $\text{IC}_{\bar{y}}$ and IC_y accounts for the missing numerator, which also corresponds to the factor of $\frac{1}{L(1, \chi, \bar{n})}$ in the Mellin transform (3.3).

4.5. **Crystals.** To reconnect with Theorem 3.7.2, define V_X^+ to be the \check{T} -representation with basis in bijection with \mathfrak{B}^+ . The crux of Conjecture 1.5.1 is getting half of a \check{G} -representation ρ_X .

Since we know this is what we want, formally set $\mathfrak{B} = \mathfrak{B}^+ \sqcup (\mathfrak{B}^+)^*$, where $(\mathfrak{B}^+)^*$ is defined to be in bijection with \mathfrak{B}^+ but the weights are replaced by their negatives. In this way, $(\mathfrak{B}^+)^*$ naturally corresponds to a basis of $(V_X^+)^*$.

Theorem 4.5.1 (Sakellaridis–W). *\mathfrak{B} has the structure of a (Kashiwara) crystal, i.e., a graph with weighted vertices and edges corresponding to lowering operators \tilde{f}_α .*

We use this abstract combinatorial notion of crystal as a bridge to hopefully getting a crystal basis. A crystal basis is the (Lusztig) canonical basis¹⁵ at $q = 0$ of an integrable $U_q(\mathfrak{g})$ -module in category \mathcal{O} . So the crystal basis is a way for us to access a \check{G} -representation.

$$\text{f.d. } \check{G}\text{-representation} \rightsquigarrow \text{crystal basis} \in \{\text{crystals}\}$$

Conjecture 4.5.2. *\mathfrak{B} is the crystal basis for a finite dimensional \check{G} -representation ρ_X .*

This conjecture implies Conjecture 1.5.1 (by construction, \mathfrak{B} corresponds to a basis of $\rho'_X = T^*V_X^+$).

4.6. **Further details.** We can identify $(\text{Gr}_B^{\check{\lambda}})_{\text{red}} = \mathbf{N}_{\mathbf{F}} t^{\check{\lambda}} \mathbf{G}_{\mathbf{O}} / \mathbf{G}_{\mathbf{O}} =: S^{\check{\lambda}} \subset \text{Gr}_G$, i.e., a semi-infinite orbit. Let $\bar{S}^{\check{\lambda}}$ denote its closure in Gr_G . Then the fiber of $\bar{y} \rightarrow \mathcal{A}$ over $\check{\lambda} \cdot v$ is

$$\bar{Y}^{\check{\lambda}} = \bar{S}^{\check{\lambda}} \times_{\mathbf{X}_{\mathbf{F}}/\mathbf{G}_{\mathbf{O}}} \mathbf{X}_{\mathbf{O}}/\mathbf{G}_{\mathbf{O}}.$$

Proposition 4.6.1 ([MV]). *The boundary $\bar{S}^{\check{\lambda}} = \bigcup_{\check{\mu} \leq \check{\lambda}} S^{\check{\mu}}$ is given by a hyperplane section in Gr_G .*

¹⁵Canonical bases were first discovered by Lusztig '90 in types A, D, E, and subsequently by Kashiwara using different methods. The crystal basis at $q = 0$ in types A, B, C, D was discovered independently by Kashiwara at around the same time in '90.

We have $\bar{Y}^\lambda \cap S^\lambda = S^\lambda \times_{\mathbf{X}_F/\mathbf{G}_O} \mathbf{X}_O/\mathbf{G}_O$, which further breaks up according to \mathbf{G}_O -orbits of \mathbf{X}_O into pieces

$$(\mathbf{N}_F t^\lambda \mathbf{G}_O \cap H_F t^\theta \mathbf{G}_O)/\mathbf{G}_O$$

with $\theta \in -\check{\Lambda}_G^+$.

The lowering operator we define on \mathfrak{B} is roughly given by

$$\bar{Y}^\lambda \rightsquigarrow \bar{Y}^\lambda \cap S^{\lambda-\check{\alpha}} \subset Y^{\lambda-\check{\alpha}}.$$

This does not quite uniquely specify how to lower an irreducible component to another irreducible component, but a reduction to considering affine embeddings of $\mathbb{G}_m \backslash \mathrm{GL}_2 \times (\text{torus})$ gives us enough information to pick out the correct irreducible component in $\bar{Y}^{\lambda-\check{\alpha}}$.

For a summary of the proofs of our main results, I refer to [SW, §1.3-1.5] of our paper.

REFERENCES

- [BBDG] A. A. Beilinson, J. Bernstein, P. Deligne, and O. Gabber, *Faisceaux pervers*, Analysis and topology on singular spaces, I (Luminy, 1981), Astérisque, vol. 100, Soc. Math. France, Paris, 2018, pp. 5171.
- [BZSV] D. Ben-Zvi, Y. Sakellaridis, A. Venkatesh, Relative duality, in preparation.
- [Ber] J. Bernstein, *On the support of Plancherel measure*.
- [BP] R. Beuzart-Plessis, Comparison of local spherical characters and the Ichino-Ikeda conjecture for unitary groups, Journal of the Institute of Mathematics of Jussieu, 1-52. doi:10.1017/S1474748019000707
- [BK] R. Bezrukavnikov and D. Kazhdan, *Geometry of second adjointness for p -adic groups*, Represent. Theory 19 (2015), 299332, With an appendix by Yakov Varshavsky, Bezrukavnikov and Kazhdan.
- [BNS] A. Bouthier, B. C. Ngô, and Y. Sakellaridis, *On the formal arc space of a reductive monoid*, Amer. J. Math. **138** (2016), no. 1, 81–108.
- [BFGM] A. Braverman, M. Finkelberg, D. Gaitsgory, and I. Mirković, *Intersection cohomology of Drinfeld's compactifications*, Selecta Math. (N.S.) **8** (2002), no. 3, 381–418.
- [BrK] A. Braverman and D. Kazhdan, γ -functions of representations and lifting. With an appendix by V. Vologodsky. GAFA 2000 (Tel Aviv, 1999). Geom. Funct. Anal. 2000, Special Volume, Part I, 237278.
- [D] V. Drinfeld, *Grinberg-Kazhdan theorem and Newton groupoids*, 2018.
- [GN] Dennis Gaitsgory and David Nadler, *Spherical varieties and Langlands duality*, Mosc. Math. J. **10** (2010), no. 1, 65–137, 271.
- [GW] Wee Teck Gan, Xiaolei Wan, *Relative character identities and theta correspondence*.
- [II] Atsushi Ichino and Tamutsu Ikeda. *On the periods of automorphic forms on special orthogonal groups and the Gross-Prasad conjecture*. Geom. Funct. Anal., 19(5):13781425, 2010.
- [J] Hervé Jacquet. Factorization of period integrals. J. Number Theory, 87(1):109–143, 2001. doi:10.1006/jnth.2000.2584.
- [KS] F. Knop and B. Schalke, *The dual group of a spherical variety*, Trans. Moscow Math. Soc. **78** (2017), 187–216.
- [KSt] F. Knop and B. Van Steirteghem, *Classification of smooth affine spherical varieties*. Transform. Groups 11 (2006), no. 3, 495516.
- [Lun73] Domingo Luna, *Slices étales*, Sur les groupes algébriques, 1973, pp. 81–105. Bull. Soc. Math. France, Paris, Mémoire 33. MR 0342523
- [MV] I. Mirković and K. Vilonen, *Geometric Langlands duality and representations of algebraic groups over commutative rings*, Ann. of Math. (2) **166** (2007), no. 1, 95–143.
- [N] B. C. Ngô, *On a certain sum of L -functions*.
- [Ric77] R. W. Richardson, *Affine coset spaces of reductive algebraic groups*, Bull. London Math. Soc. **9** (1977), no. 1, 38–41. MR 437549
- [Sa08] Y. Sakellaridis, *On the unramified spectrum of spherical varieties over p -adic fields*, Compositio Mathematica 144 (2008), no. 4, 9781016.
- [Sa12] ———, *Spherical varieties and integral representations of L -functions*, Algebra Number Theory **6** (2012), no. 4, 611–667, Updated arXiv version.
- [Sa13] ———, *Spherical functions on spherical varieties*, Amer. J. Math. **135** (2013), no. 5, 1291–1381.
- [Sa18] ———, *Inverse Satake Transforms*, Geometric Aspects of the Trace Formula (Cham), Springer International Publishing, 2018, pp. 321349.

- [SV] Y. Sakellaridis and A. Venkatesh, *Periods and harmonic analysis on spherical varieties*, *Astérisque* (2017), no. 396, viii+360.
- [SW] Y. Sakellaridis and J. Wang, *Local L -factors and geometric asymptotics for spherical varieties*, arXiv:2009.03943
- [W] J. Wang, *On an invariant bilinear form on the space of automorphic forms via asymptotics*, *Duke Math. J.* 167 (16), 2965-3057 (2018).
- [Was] B. Wasserman, *Wonderful varieties of rank two*, *Transform. Groups* 1 (1996), no. 4, 375-403.